

Carcinoma originating from aberrant breast tissue. A case report and review of the literature

Sibel Kahraman-Cetintas¹, Senem Turan-Ozdemir², Ugur Topal³, Meral Kurt¹,
Sehsuvar Gokgoz⁴, Ozlem Saraydaroglu⁵, and Lutfi Ozkan¹

¹Department of Radiation Oncology, ²Department of Anatomy, ³Department of Radiology, ⁴Department of Surgery, and ⁵Department of Pathology, Uludag University Faculty of Medicine, Bursa, Turkey

ABSTRACT

Carcinoma arising from ectopic breast tissue, either supernumerary breast or aberrant breast tissue, is extremely rare. Carcinoma occurs more frequently in the ectopic breast tissue of the axilla than in extra-axillary ectopic breast tissue. Here we report a case of an invasive lobular carcinoma arising from extra-axillary ectopic breast tissue and presenting as a subcutaneous nodule.

Key words: *ectopic breast, supernumerary breast, accessory breast, aberrant breast tissue, breast cancer.*

Correspondence to: Sibel Kahraman-Cetintas, Department of Radiation Oncology, Faculty of Medicine, Uludag University, 16059 Bursa, Turkey.

Tel +90-224-442 91 60;
e-mail skahraman@uludag.edu.tr
skahramanc@hotmail.com

Received January 10, 2007;
accepted August 29, 2007.